

Nonlinearity Management in Higher Dimensions

P.G. Kevrekidis¹, D.E. Pelinovsky², and A. Stefanov³

¹ Department of Mathematics and Statistics, University of Massachusetts, Amherst, MA 01003

² Department of Mathematics, McMaster University, Hamilton, Ontario, Canada, L8S 4K1

³ Department of Mathematics, University of Kansas, 1460 Jayhawk Blvd, Lawrence, KS 66045–7523

September 13, 2005

Abstract

In the present short communication, we revisit nonlinearity management of the time-periodic nonlinear Schrödinger equation and the related averaging procedure. By means of rigorous estimates, we show that the averaged nonlinear Schrödinger equation does not blow up in the higher dimensional case so long as the corresponding solution remains smooth. In particular, we show that the H^1 norm remains bounded, in contrast with the usual blowup mechanism for the focusing Schrödinger equation. This conclusion agrees with earlier works in the case of strong nonlinearity management but contradicts those in the case of weak nonlinearity management. The apparent discrepancy is explained by the divergence of the averaging procedure in the limit of weak nonlinearity management.

1 Introduction

In the past few years, there has been a large volume of literature regarding applications of the nonlinear Schrödinger equation (NLS) in the presence of the so-called *nonlinearity management* (often referred to also as Feshbach resonance management). The NLS is a prototypical dispersive nonlinear wave equation of the form:

$$iu_t = -\Delta u + \Gamma(t)|u|^2u + V(x)u, \quad (1)$$

where $u(x, t)$ is a complex envelope field, $V(x) \geq 0$ is an external potential, $\Gamma(t)$ is a time-periodic nonlinearity coefficient, and Δ is the Laplacian operator with $x \in \mathbb{R}^d$, $d \geq 1$.

Nonlinearity management arises in optics for transverse beam propagation in layered optical media [1], as well as in atomic physics for the Feshbach resonance of the scattering length of inter-atomic interactions in Bose-Einstein condensates (BECs) [2]. In the latter case, the periodic variation of $\Gamma(t)$ through an external magnetic field has been used as a means of producing robust matter-wave breathers in quasi-one-dimensional BECs [3]. It has also been suggested that the nonlinearity management may prevent collapse-type phenomena in higher dimensions [4]. Theoretical studies of the collapse-type behavior were performed with a reduction of the time-periodic PDE problem (1) to a time-periodic ODE problem using the variational method [5] and the method of moments [6]. Recent work [7] presented a rigorous proof that the NLS equation (1) with $\Gamma(t) < 0$ for all $t \geq 0$ may have a blow up in three dimensions ($d = 3$) depending on parameters of the initial condition.

The physical relevance of the time-periodic NLS equation, as evidenced by the above works, led to further developments in analysis of the evolution problem (1). As an example, the success of the averaging theory [8] for optical solitons in the presence of *strong dispersion management*, led to an analogous development for *strong nonlinearity management* produced originally in [9] and systematized in [10]. Let ϵ be a small parameter that measures a short period and large variations of $\Gamma(t)$ in the form

$$\Gamma = \gamma_0 + \frac{1}{\epsilon} \gamma \left(\frac{t}{\epsilon} \right), \quad (2)$$

where $\gamma(\tau)$, $\tau = t/\epsilon$ has a unit period and zero mean. The time-periodic NLS equation (1) is averaged in the limit $\epsilon \rightarrow 0$. The averaging procedure is based on the transformation of solutions of the NLS equation (1),

$$u = e^{i\gamma_{-1}(\tau)|v|^2} v(x, t),$$

where $\gamma_{-1}(\tau)$ is the mean-zero anti-derivative of $\gamma(\tau)$, and subsequent expansion of $v(x, t)$ into an asymptotic series,

$$v = w(x, t) + \epsilon v_1(x, \tau, t) + O(\epsilon^2).$$

The secular growth of $v_1(x, \tau, t)$ in τ is removed if $w(x, t)$ satisfies the averaged NLS equation,

$$iw_t = -\Delta w + \gamma_0 |w|^2 w + V(x)w - \sigma^2 \left(|\nabla |w|^2|^2 + 2|w|^2 \Delta |w|^2 \right) w, \quad (3)$$

where

$$\sigma^2 = \int_0^1 \gamma_{-1}^2(\tau) d\tau.$$

Our paper addresses the question whether the averaged NLS equation (3) with $\gamma_0 < 0$ and $\sigma \neq 0$ arrests the blowup of solutions of the NLS equation in two and three dimensions. We remind to readers that the averaged NLS equation (3) with $\gamma_0 < 0$ and $\sigma = 0$ may have blowup in $d \geq 2$ depending on the initial condition [11]. A similar question was addressed recently in [12] in the context of the cubic-quintic NLS equation in one dimension ($d = 1$).

Before we explain our results in more detail, let us take a moment to review the usual blow up mechanism in the context of a semilinear focusing Schrödinger equation. That is, take $V(x) = 0$ and $\sigma = 0$ in the NLS equation (3). It is well-known (see Chapter 5 in [13]) that whenever $d \geq 2$ and $\gamma_0 < 0$, and for generic (smooth) initial data $u_0(x)$ in H^1 , there exists a time $T^* < \infty$, so that

$$\lim_{t \rightarrow T^*} \sup_{0 < t < T^*} \int_{\mathbb{R}^d} |\nabla u(x, t)|^2 dx \rightarrow \infty \quad \text{and} \quad \lim_{t \rightarrow T^*} \sup_{0 < t < T^*} \int_{\mathbb{R}^d} |u(x, t)|^4 dx \rightarrow \infty. \quad (4)$$

while the solution is sufficiently smooth in $t \in (0, T^*)$, and thus for $0 \leq t < T^*$,

$$H[u(\cdot, t)] = \int_{\mathbb{R}^d} \left(\frac{1}{2} |\nabla u(x, t)|^2 + \frac{\gamma_0}{4} |u(x, t)|^4 \right) dx = H[u_0(\cdot)].$$

In our work, we show that no such blowup of solutions occurs in the averaged NLS equation (3) with $\gamma_0 < 0$ and $\sigma \neq 0$. That is, assuming that the solution exists and satisfies the conservation law $H[u(\cdot, t)] = H[u_0]$ and $P[u] = \|u(\cdot, t)\|_{L^2}^2 = \|u_0\|_{L^2}^2$, one can in fact control *a priori* $\|u(\cdot, t)\|_{H^1}$ and $\|u(\cdot, t)\|_{L^4}$.

Unfortunately, our arguments do not allow to rigorously justify the global (and in fact even the local) existence of solutions, satisfying the physical conservation laws. In other words, the solutions may still blow up due to eventual loss of smoothness (and a subsequent failure to obey the conservation laws), although we consider that to be an unlikely scenario, especially for sufficiently smooth initial data $u_0(x)$.

We demonstrate our claims initially, in Section 2, from the point of view of scaling arguments. Then using rigorous estimates in Section 3, we show that the H^1 and the L^4 norm of the solution $u(x, t)$ are controlled by $H[u_0]$ and $P[u_0]$. In Section 4 we compare the above conclusion and earlier works where blowup of solutions of the full time-periodic NLS equation (1) have been reported. Section 5 summarizes our findings.

2 Formal Scaling Arguments

The averaged NLS equation (3) has a standard Hamiltonian form (see [10]) with the Hamiltonian functional:

$$H(w) = \int_{\mathbb{R}^d} \left(|\nabla w|^2 + \frac{\gamma_0}{2} |w|^4 + V(x) |w|^2 + \sigma^2 |w|^2 |\nabla |w|^2|^2 \right) dx. \quad (5)$$

Due to the gauge invariance, the averaged NLS equation (3) also conserves the squared L^2 norm:

$$P(w) = \int_{\mathbb{R}^d} |w|^2 dx. \quad (6)$$

Solitary wave solutions of the averaged NLS equation (3) are critical points of $H(w)$ at the level set of fixed values of $P(w)$.

Using formal scaling arguments [14] (see also [11]), we consider a two-parameter family of dilatations:

$$w = bW(ax), \quad (7)$$

where (a, b) are parameters and $W(\xi)$ is a suitable function of $\xi = ax$. The squared L^2 norm (6) is preserved by the dilatations (7) whenever $b = a^{d/2}$. The Hamiltonian (5) at the dilatations (7) is scaled as a function of parameter $a > 0$:

$$H(a) = I_0(a) + a^2 I_1 + \gamma_0 a^d I_2 + \sigma^2 a^{2d+2} I_3, \quad (8)$$

where

$$I_1 = \int_{\mathbb{R}^d} |\nabla W|^2 d\xi, \quad I_2 = \frac{1}{2} \int_{\mathbb{R}^d} |W|^4 d\xi, \quad I_3 = \int_{\mathbb{R}^d} |W|^2 |\nabla |W|^2|^2 d\xi.$$

and

$$I_0(a) = \int_{\mathbb{R}^d} V\left(\frac{\xi}{a}\right) |W|^2 d\xi.$$

Let us consider the case of no nonlinearity management and no external potential, when $\sigma^2 = 0$ and $V(x) = 0$. It follows from (8) that the Hamiltonian function $H(a)$ is positive definite in the defocusing case, when $\gamma_0 > 0$. In the focusing case, when $\gamma_0 < 0$, the Hamiltonian function $H(a)$ is bounded from below for $d = 1$ and $d = 2$, $\gamma_{\text{cr}} < \gamma_0 < 0$ and is unbounded from below for $d = 2$, $\gamma_0 < \gamma_{\text{cr}}$ and $d \geq 3$, where

$$\gamma_{\text{cr}} = -\frac{I_1}{I_2}.$$

When $H(a)$ is unbounded from below as $a \rightarrow \infty$, the critical points of $H(w)$ at a fixed value of $P(w)$ (i.e., solitary wave solutions) can not be stable for small width a^{-1} and instability of solitary waves implies a blowup of localized initial data in the time evolution of the cubic NLS equation (see [11] for details).

When the nonlinearity management is applied, the last term in the decomposition (8) always dominates and it preserves the boundness of $H(a)$ from below for any $\sigma^2 > 0$. This indicates on the level of formal scaling arguments that the blowup of solutions is arrested by the nonlinearity management term in the averaged NLS equation (3).

We note that the first term in the decomposition (8) does not change the conclusions above if $V(x)$ is a smooth non-negative potential, such that $I_0(a) \geq 0$. Typical examples of $V(x)$ are parabolic magnetic traps, when $V \sim x^2$, and periodic optical lattices, when $V \sim \sin^2(x)$.

3 On Global Solutions to the Averaged Equation

The rigorous analysis of local well-posedness of the averaged NLS equation (3) is not a trivial task. In fact, to the best of our knowledge, one cannot verify even the local existence and uniqueness of solutions to this problem. The problem has been considered in one dimension $d = 1$ by Poppenberg where existence and uniqueness of local solutions with initial data in H^∞ was established [15]. In higher dimensions $d \geq 2$, one needs to require [16] that the initial data be in $H^{s,m}$ for sufficiently large values of s, m , where

$$H^{s,m} = \left\{ f \in C^s(\mathbb{R}^d; \mathbb{C}) : \int_{\mathbb{R}^d} (1 + |x|)^{2m} (|f|^2 + |\partial^s f|^2) dx < \infty \right\}.$$

In addition, one needs to assume a “non-trapping” geometric condition on the symbol of the second order operator, which depends on the profile of the initial data (see [16] for details).

On the contrary, the local well-posedness of the time-periodic NLS equation (1) can be proved with the standard tool in the energy space $H^1 \equiv H^{1,0}$ (see Section 2 in [17]). Therefore, one is tempted to assume that solutions of the averaged NLS equation (3), derived from the time-periodic NLS equation (1) with a regular asymptotic procedure, inherit local well-posedness. However, we cannot state a precise condition under which the averaged NLS equation (3) has a (local) solution that conserves constant values of P and H in time.

Thus, as we have explained in Section 1, we will focus our attention on the following problem: Assuming the existence of a local solution in space

$H^1 \cap L^4 \cap \{w : \int |w|^2 |\nabla w|^2 dx < \infty\}$ and in a time interval $t \in (0, T)$, we will show that

$$\sup_{0 < t < T} (\|\nabla u(\cdot, t)\|_{L^2} + \|u\|_{L^4}) < \infty.$$

In other words, the standard blow up mechanism (4) for the cubic focusing Schrödinger equation does not occur.

To that end, we represent the Hamiltonian $H(w)$ in the form:

$$H(w) = H_1(w) + \gamma_0 H_2(w), \quad (9)$$

where

$$H_1(w) = \int_{\mathbb{R}^d} (|\nabla w|^2 + V(x)|w|^2 + \sigma^2 |w|^2 |\nabla w|^2) dx \geq 0$$

and

$$H_2(w) = \frac{1}{2} \int_{\mathbb{R}^d} |w|^4 dx \geq 0.$$

We consider the focusing case $\gamma_0 < 0$ and prove that $H_1(w)$ and $H_2(w)$ are bounded by the two conserved quantities $H(w)$ and $P(w)$.

First, we quote a variant of the Gagliardo-Nirenberg inequality (see (1.1.16) on p. 15 in [18]).

Lemma 1. *For all $1 \leq p, q, r \leq \infty$, $\theta \in (0, 1)$ and $r^{-1} = \theta p^{-1} + (1 - \theta)q^{-1}$, it is true for every function $f(x)$ on $x \in \mathbb{R}^d$ that*

$$\|f\|_{L^r} \leq \|f\|_{L^p}^\theta \|f\|_{L^q}^{1-\theta}. \quad (10)$$

Next, we quote a modification of the Sobolev embedding theorem (see (19) on p.26 in [19]).

Lemma 2. *There exists a constant $C_d > 0$, which depends only on the dimension $d \geq 1$, so that it is true for every function $f(x)$ on $x \in \mathbb{R}^d$ that*

$$\|f\|_{L^2} \leq C_d (\|\nabla f\|_{L^2} + \|f\|_{L^1}). \quad (11)$$

The main result of our analysis is the following theorem.

Theorem 1. *There exist $\mu(d) > 0$ and $C(\mu, d, \sigma) > 0$, so that for every $0 < \mu < \mu(d)$ and every function $\phi(x)$ on $x \in \mathbb{R}^d$, it is true that*

$$\|\phi\|_{L^4}^4 \leq \mu H_1(\phi) + C(\mu, d, \sigma) \left(\|\phi\|_{L^2}^2 + \|\phi\|_{L^2}^4 \right). \quad (12)$$

Proof. Let $f = h^{3/2}$ with $h(x) > 0$ in (11) and obtain

$$\int_{\mathbb{R}^d} h^3 dx \leq C_d^2 \left[\frac{3}{2} \left(\int_{\mathbb{R}^d} h |\nabla h|^2 dx \right)^{1/2} + \int_{\mathbb{R}^d} h^{3/2} dx \right]^2.$$

Next, we set $h = |\phi|^2$ and obtain

$$\begin{aligned} \int_{\mathbb{R}^d} |\phi|^6 dx &\leq C_d^2 \left[\frac{3}{2} \left(\int_{\mathbb{R}^d} |\phi|^2 |\nabla |\phi|^2|^2 dx \right)^{1/2} + \int_{\mathbb{R}^d} |\phi|^3 dx \right]^2 \\ &\leq C_d^2 \left[\frac{9}{2} \int_{\mathbb{R}^d} |\phi|^2 |\nabla |\phi|^2|^2 dx + 2 \left(\int_{\mathbb{R}^d} |\phi|^3 dx \right)^2 \right] \\ &\leq C_{d,\sigma} \left[H_1(\phi) + \|\phi\|_{L^3}^6 \right], \end{aligned}$$

for some positive constant $C_{d,\sigma}$. We have used here that

$$(\sqrt{a} + b)^2 \leq 2(a + b^2).$$

By the Gagliardo-Nirenberg inequality (10), we have $\|\phi\|_{L^3} \leq \|\phi\|_{L^2}^{1/2} \|\phi\|_{L^6}^{1/2}$, such that the last inequality is rewritten in the form:

$$\begin{aligned} \|\phi\|_{L^6}^6 &\leq C_{d,\sigma} H_1(\phi) + C_d^2 \|\phi\|_{L^6}^3 \|\phi\|_{L^2}^3 \\ &\leq C_{d,\sigma} H_1(\phi) + C_d^2 \left[\mu \|\phi\|_{L^6}^6 + \frac{1}{4\mu} \|\phi\|_{L^2}^6 \right], \end{aligned} \quad (13)$$

where we have used the Cauchy-Schwartz inequality:

$$\forall \mu > 0 : ab \leq \mu a^2 + \frac{b^2}{4\mu}. \quad (14)$$

Let $\mu < \mu(d)$, where $2C_d^2\mu(d) = 1$. The term $\|\phi\|_{L^6}^6$ can be estimated from the bound (13) as follows:

$$\|\phi\|_{L^6}^6 \leq \tilde{C}_{d,\sigma} H_1(\phi) + \tilde{C}_d \|\phi\|_{L^2}^6 \quad (15)$$

for some constants $\tilde{C}_{d,\sigma} > 0$ and $\tilde{C}_d > 0$. By the Gagliardo-Nirenberg inequality (10), we have $\|\phi\|_{L^4} \leq \|\phi\|_{L^6}^{3/4} \|\phi\|_{L^2}^{1/4}$, such that the upper bound for $\|\phi\|_{L^4}$ follows from (14) and (15):

$$\begin{aligned} \|\phi\|_{L^4}^4 &\leq \|\phi\|_{L^2} \left[\hat{C}_{d,\sigma} \sqrt{H_1(\phi)} + \hat{C}_d \|\phi\|_{L^2}^3 \right] \\ &\leq \frac{\hat{C}_{d,\sigma}}{4\mu} \|\phi\|_{L^2}^2 + \hat{C}_d \|\phi\|_{L^2}^4 + \mu H_1(\phi), \end{aligned}$$

which is the desired upper bound (12). \square

As a corollary of the main theorem, we pick $\mu = \mu(d)/2$ and immediately obtain the following bounds on the two parts of the energy functionals (9).

Corollary 1. *There exists constants $C_1 > 0$ and $C_2 > 0$ that depend on d, γ_0, σ , so that*

$$H_1(w) \leq C_1 (H(w) + P(w) + P^2(w)) \quad (16)$$

and

$$H_2(w) \leq C_2 (H(w) + P(w) + P^2(w)). \quad (17)$$

Thus, if $H(w)$ and $P(w)$ are conserved by the time evolution, the Cauchy problem for the averaged NLS equation (3) has global solutions in H^1 . Therefore, assuming that physical relevant solutions conserve Hamiltonian H and charge P , we conclude that the blowup of solutions of the cubic NLS equation (3) with $\gamma_0 < 0$ in $d \geq 2$ is arrested by the nonlinearity management for any $\sigma \neq 0$.

4 Averaged Equation versus Full Dynamics

As we have shown in Section 3, solutions of the averaged NLS equation (3) with $\gamma_0 < 0$ and $\sigma \neq 0$, which preserve $H(w)$ and $P(w)$, do not blowup in higher dimensions. This result raises the question whether the blowup of solutions is arrested in the time-periodic NLS equation (1) for any non-zero variance of the nonlinearity coefficient $\Gamma(t)$. We address this question within the ODE reduction of the time-periodic problem, which was considered recently through the variational method [4, 5] and the method of moments [6]. We note that the ODE reductions imply that the solution $u(x, t)$ remains infinitely smooth and the blowup may only occur due to infinite increase in certain norms. Therefore, the blowup in the ODE approach would correspond directly to the blowup estimates obtained in Section 3.

It follows from the method of moments [6] that the time evolution of the radially symmetric localized solutions of the full NLS equation (1) is approximated by a time-dependent, generalized Ermakov-Pinney [20] equation:

$$\ddot{R}(t) = \frac{Q_1}{R^3} + \Gamma(t) \frac{Q_2}{R^{d+1}}, \quad (18)$$

where $R(t) \geq 0$ is an effective width of a localized solution, while (Q_1, Q_2) are constants found from an initial data, such that $Q_2 > 0$ (see Section 3 in [6] for details). We shall consider the critical case $d = 2$ and rewrite the ODE (18) with the nonlinearity coefficient $\Gamma(t)$ in (2) in the explicit form:

$$\ddot{R}(t) = \frac{\alpha + \beta \gamma(t/\epsilon)}{R^3}, \quad (19)$$

where $\alpha = Q_1 + \gamma_0 Q_2$, $\beta = Q_2/\epsilon > 0$, and $\gamma(\tau)$, $\tau = t/\epsilon$ is a periodic function with zero mean and unit period. Conditions for blowup and existence of bounded oscillations in solutions of the ODE (19) were found in [21] (see Section 4 in [6]). The sufficient condition for the blowup (when $R(t) \rightarrow 0$ in a finite time $t \rightarrow t_0$) is

$$\alpha + \beta \max_{0 \leq \tau \leq 1} (\gamma) < 0. \quad (20)$$

The necessary condition for the bounded oscillations of $R(t) > 0$ for any $t \geq 0$ is

$$\alpha + \beta \max_{0 \leq \tau \leq 1} (\gamma) > 0. \quad (21)$$

Numerical simulations in the case $\alpha < 0$ showed that the condition (21) was also sufficient for the blowup arrest for any $t \geq 0$ [6].

It is obvious from the definition of α and β that $\beta \gg |\alpha|$ in the asymptotic limit $\epsilon \rightarrow 0$. Therefore, the condition (21) is satisfied and solutions of the time-periodic ODE (19) do

not collapse, in agreement with our results derived in the context of the averaged NLS equation (3).

Previous work [4] (see also the review in [10]) has also addressed the averaged NLS equation (3) in the limit of *weak* nonlinearity management, when $\gamma(\tau)$ and σ^2 are rescaled as $\gamma = \epsilon\tilde{\gamma}(\tau)$ and $\sigma^2 = \epsilon^2\tilde{\sigma}^2$ such that σ^2 is small in the limit $\epsilon \rightarrow 0$. Solutions of the averaged NLS equation (3) with $\sigma^2 = \epsilon^2\tilde{\sigma}^2$ blow up in a finite time for $\epsilon = 0$, but the small ϵ^2 -terms formally stabilize the blowup for any $\tilde{\sigma}^2 > 0$. Although the truncation of the critical NLS equation by neglecting terms of the order of ϵ^4 has been considered in many applications of nonlinear optics (see Sections 4-5 in [22] and references therein), it is clearly insufficient for a correct identification of the domain, where the blowup of solutions occurs. Indeed, while the averaged NLS equation (3) with small σ^2 predicts no blowup of solutions, the full NLS equation (1) with $\Gamma(t) < 0$ may have the blow up, according to recent results [7].

The weak nonlinearity management corresponds to the case $\beta \approx |\alpha|$. Both domains (20) and (21) for blowup and bounded oscillations fit to the limit of weak nonlinearity management. However, we will show that the averaged equation can not distinguish between these two different cases of global dynamics in solutions of the ODE problem (19). Following [5], we consider a simple time-periodic ODE:

$$\ddot{R}(t) = \frac{\alpha + \beta \sin(2\pi\tau)}{R^3}, \quad \tau = \frac{t}{\epsilon}, \quad (22)$$

where $\alpha < 0$, $\beta > 0$ and (α, β) are order of $O(1)$ in the limit $\epsilon \rightarrow 0$. By using the formal asymptotic multi-scale expansion method (see [10] for details), we construct an asymptotic solution to the problem (22):

$$R = r(t) + \epsilon^2 R_2(\tau, r) + \epsilon^4 R_4(\tau, r) + O(\epsilon^6), \quad \tau = \frac{t}{\epsilon}, \quad (23)$$

where R_2 and R_4 are recursively found from the set of linear inhomogeneous problems,

$$\begin{aligned} R_2 &= -\frac{\beta}{(2\pi)^2 r^3} \sin(2\pi\tau), \\ R_4 &= -\frac{3\alpha\beta}{(2\pi)^4 r^7} \sin(2\pi\tau) + \frac{3\beta^2}{8(2\pi)^4 r^7} \cos(4\pi\tau). \end{aligned}$$

The mean-value term $r(t)$ satisfies an extended dynamical equation that excludes secular growth of the correction terms of the series (23) in τ :

$$\ddot{r} = \frac{\alpha}{r^3} + \epsilon^2 \frac{3\beta^2}{2r^7} + \epsilon^4 \frac{15\alpha\beta^2}{2r^{11}} + O(\epsilon^6), \quad \epsilon = \frac{\epsilon}{2\pi}. \quad (24)$$

The averaged ODE problem (24) is an equation of motion for an effective particle with a coordinate $r(t)$ in the potential field with an effective potential energy:

$$U(r) = \frac{\alpha}{2r^2} + \epsilon^2 \frac{\beta^2}{4r^6} + \epsilon^4 \frac{3\alpha\beta^2}{4r^{10}} + O(\epsilon^6). \quad (25)$$

When $\alpha < 0$ and $\epsilon = 0$, the particle with $r(0) > 0$ reaches $r = 0$ in a finite time $t = t_0 < \infty$, that indicates the blowup of a localized solution. When the next ϵ^2 -term is taken into

account (as in the approximation of weak nonlinearity management [4, 5]), the blowup is arrested and the mean-value term $r(t)$ oscillates in an effective minimum of the potential energy $U(r)$, truncated at ε^2 -terms. When the next ε^4 -term is taken into account (beyond the approximation of weak nonlinearity management), the potential energy $U(r)$ with $\alpha < 0$ does not prevent the blowup of the localized solution depending on the initial data $r(0)$. Existence versus non-existence of blowup depends on the ratio of parameters (α, β) but the difference is detected within the averaging method only if convergence of the power series (25) is established in a closed analytical form.

Similarly, the averaged NLS equation (3) can not be used in the limit of weak nonlinearity management for an accurate prediction of existence versus non-existence of blowup of solutions. In order to illustrate this point, we have performed numerical simulations of the time-periodic NLS equation (1) in two dimensions ($d = 2$) with

$$\Gamma(t) = -20 + 8 \sin(2\pi t),$$

where numerical values in $\Gamma(t)$ correspond to the sufficient condition for blowup (20) to occur. We have observed numerically that collapse of radially symmetric Gaussian initial data does occur (see solid curve on Fig. 1) by monitoring the width of the one-dimensional slice along $y = 0$ of the wavefunction,

$$\bar{x} = \left(\frac{\int x^2 |u(x, 0, t)|^2 dx}{\int |u(x, 0, t)|^2 dx} \right)^{1/2},$$

until it becomes comparable to the lattice grid spacing used (at that scale collapse is arrested, since the numerical scheme cannot resolve scales below the grid spacing). On the other hand, numerical simulations of the averaged NLS equation (3) with the same parameters show that the width \bar{x} never decreased below $\bar{x} < 0.45$ (see dashed curve on Fig. 1) indicating the absence of collapse in accordance with results of Section 3. Similar outcomes of numerical simulations were observed for other numerical values in $\Gamma(t)$, as soon as the sufficient condition for blowup (20) was satisfied.

We summarize that the averaged NLS equation (3) can only be used for modeling of the blowup arrest in the limit of strong nonlinearity management of the full NLS equation (1). The averaged NLS equation can not distinguish between the blowup and no blowup domains in the limit of weak nonlinearity management.

5 Conclusion

In conclusion, we have studied the question whether solutions of the averaged NLS equation blow up in finite time. The averaged NLS equation describes strong nonlinearity management of the time-periodic NLS equation. We have showed with formal scaling arguments and rigorous analysis that the blowup of solutions in higher dimensions is arrested within the averaged NLS equation. We have also discussed the non-applicability of the averaged NLS equation to the weak nonlinearity management, where the blowup of solutions can occur beyond the weak management limit.

It is an open problem to study conditions for blowup in the time-periodic NLS equation, depending on parameters of the nonlinearity management and profile of initial data.




Figure 1: Numerical simulations of the full NLS equation (1) (solid curve) and the averaged NLS equation (3) with a fourth order in time scheme, where spacings are $dx = 0.075$ and $dt = 10^{-5}$. The half-width of the wavefunction $\bar{x}/2$ is shown as a function of time t .

Rigorous results on the latter problem are only available within the ODE approximation (18), when the PDE problem reduces to a dynamical system with one degree of freedom. It would be particularly interesting to study mathematically and to examine numerically whether the theoretical prediction from the method of moments provides an optimal bound for the full PDE problem with arbitrary initial data.

Acknowledgements. The first author is supported in part by NSF-DMS 0204585, NSF-DMS-0505063, an NSF CAREER award and the Eppley Foundation for Research. The second author is supported by NSERC and PREA grants. The third author supported in part by NSF-DMS 0300511.

The authors are thankful to Prof. D.E. Frantzeskakis for numerous enlightening discussions on the subject. We also would like to thank Prof. E. Kirr for a number of useful comments on the manuscript.

References

- [1] I. Towers and B.A. Malomed, J. Opt. Soc. Am. **19**, 537 (2002); L. Bergé et al., Opt. Lett. **25**, 1037 (2000).
- [2] S. Inouye *et al.*, Nature **392**, 151 (1998); J. Stenger *et al.*, Phys. Rev. Lett. **82**, 2422 (1999); J.L. Roberts *et al.*, Phys. Rev. Lett. **81**, 5109 (1998); S.L. Cornish *et al.*, Phys. Rev. Lett. **85**, 1795 (2000); E.A. Donley *et al.*, Nature **412**, 295 (2001).
- [3] P.G. Kevrekidis, G. Theocharis, D.J. Frantzeskakis, and B.A. Malomed, Phys. Rev. Lett. **90**, 230401 (2003).

- [4] F.Kh. Abdullaev, J.G. Caputo, R.A. Kraenkel, and B.A. Malomed, *Phys. Rev. A* **67**, 013605 (2003).
- [5] H. Saito and M. Ueda, *Phys. Rev. Lett.* **90**, 040403 (2003).
- [6] G.D. Montesinos, V.M. Pérez-García and P.J. Torres, *Physica D* **191**, 193 (2004).
- [7] V.V. Konotop and P. Pacciani, *Phys. Rev. Lett.* **94**, 240405 (2005).
- [8] I. Gabbitov and S. Turitsyn, *Opt. Lett.* **21**, 327 (1996); M.J. Ablowitz and G. Biondini, *Opt. Lett.* **23**, 1668 (1998).
- [9] D.E. Pelinovsky, P.G. Kevrekidis and D.J. Frantzeskakis, *Phys. Rev. Lett.* **91**, 240201 (2003); D.E. Pelinovsky, P.G. Kevrekidis, D.J. Frantzeskakis and V. Zharnitsky, *Phys. Rev. E* **70**, 047604 (2004).
- [10] D.E. Pelinovsky and V. Zharnitsky, *Chaos*, to appear (2006).
- [11] E.A. Kuznetsov, A.M. Rubenchik, and V.E. Zakharov, *Phys. Rep.* **142**, 103 (1986).
- [12] F.Kh. Abdullaev and J. Garnier, preprint arXiv: cond-mat/0507042.
- [13] C. Sulem, P. Sulem, *The nonlinear Schrödinger equation, self-focusing and wave collapse* (Springer–Verlag, New York, 1999).
- [14] G.H. Derrick, *J. Math. Phys.* **5**, 1252 (1964).
- [15] M. Poppenberg, *Nonlinear Anal.* **45**, 723 (2001).
- [16] C. E. Kenig, *The Cauchy problem for the quasilinear Schrödinger equation*, IAS/Park city Mathematics Series, 2002.
- [17] E. Kirr, S. Cuccagna, and D. Pelinovsky, *J. Diff. Eqs.*, to appear (2006).
- [18] L. Grafakos, *Classical and Modern Analysis*, Pearson/Prentice Hall, 2004.
- [19] J.J. Garcia–Ripoll, V.V. Konotop, B. Malomed, and V.M. Perez-Garcia, *Math. Comp. Simulat.* **62**, 21 (2003)
- [20] V.P. Ermakov, *Univ. Izv. Kiev* **20**, 1 (1880); E. Pinney, *Proc. Amer. Math. Soc.* **1**, 681 (1950). For a recent review, see e.g. P.B. Espinoza, math-ph/0002005.
- [21] C. King and A. Lesniewski, *Lett. Math. Phys.* **39**, 367 (1997).
- [22] G. Fibich and G. Papanicolaou, *SIAM J. Appl. Math.* **60**, 183 (2000).